| Variable  | Estimation | Base population                                |
|---|------------|--|
| Currently using a modern method   | Proportion | All women age 15–49<br>Married women age 15–49 |
| Currently using a traditional method  | Proportion | All women age 15–49<br>Married women age 15–49 |
| Currently using any contraceptive method  | Proportion | All women age 15–49<br>Married women age 15–49 |
| Currently using injectables   | Proportion | All women age 15–49<br>Married women age 15–49 |
| Currently using male condoms  | Proportion | All women age 15–49<br>Married women age 15–49 |
| Currently using implants  | Proportion | All women age 15–49<br>Married women age 15–49 |
| Chose method by self or jointly in past 12 months                                       | Proportion | All women age 15–49<br>Married women age 15–49 |
| Paid fees for family planning services in past 12 months                                | Proportion | All women age 15–49<br>Married women age 15–49 |
| Informed by provider about other methods  | Proportion | All women age 15–49<br>Married women age 15–49 |
| Informed by provider about side effects   | Proportion | All women age 15–49<br>Married women age 15–49 |
| Satisfied with provider: Would return and refer friend/relative to provider             | Proportion | All women age 15–49<br>Married women age 15–49 |
| Visited by health worker who talked about family planning information in past 12 months | Proportion | All women age 15–49<br>Married women age 15–49 |

## List of indicators for sampling errors

