PMA
PERFORMANCE MONITORING FOR ACTION

JOHNS HOPKINS
BLOOMBERG SCHOOL
of PUBLIC HEALTH

BILL & MELINDA GATES INSTITUTE for
POPULATION and REPRODUCTIVE HEALTH

Democratic Republic of Congo (Kinshasa)
Snapshot of Indicators
Phase 2 (2021) Cross-sectional Survey

PMA www.pmadata.org   f @PMA4Action   ⊙ @pma4action   🐦 @PMA4Action

Background Characteristics of Respondents

Distribution of women age 15 to 49 by background characteristics.

Background Characteristics	Percentage of Sample	Weighted Number	Unweighted Number
Total			
All	100.0	2369	2369
Age Group			
15-19	23.2	549	526
20-24	20.5	485	496
25-29	15.2	359	383
30-34	13.3	316	320
35-39	10.8	255	258
40-44	10.3	243	230
45-49	6.8	162	156
Marital Status			
Married, or in union	43.4	1029	1044
Unmarried, sexually active	19.8	470	472
Parity			
0-1 children	15.6	370	368
2-3 children	22.9	543	535
4+ children	20.2	477	487
Education			
Never	0.5	12	14
Primary	6.2	146	153
Secondary	75.1	1779	1779
Higher	18.2	432	423
Wealth Quintile			
Lowest	16.6	394	459
Lower	20.2	478	519
Middle	18.6	441	458
Higher	21.2	502	459
Highest	23.4	554	474

Current Use of Contraception

Percentage of women ages 15 to 49 currently using a contraceptive method, by type and background characteristics.

Background Characteristics	Any Method		Modern Method [1]		Traditional Method	
	All Women	Married [2] Women	All Women	Married [2] Women	All Women	Married [2] Women
Total						
All	45.3	54.9	27.3	30.3	18.0	24.6
Age Group						
15-19	22.4	--	16.0	--	6.3	--
20-24	55.8	57.1	37.2	34.4	18.6	22.7
25-29	60.5	61.4	37.7	38.0	22.9	23.4
30-34	51.5	55.4	33.6	37.3	17.9	18.1
35-39	55.2	64.1	28.5	30.8	26.7	33.3
40-44	48.3	51.7	20.5	21.4	27.8	30.2
45-49	25.2	31.8	8.8	12.6	16.4	19.2
Marital Status						
Married, or in union	54.9	na	30.3	na	24.6	na
Unmarried, sexually active	78.8	na	50.4	na	28.4	na
Parity						
0-1 children	53.1	48.1	30.8	21.5	22.4	26.6
2-3 children	55.5	56.5	33.6	32.6	21.9	23.9
4+ children	55.5	61.2	30.5	33.7	25.0	27.5
Education						
Never	--	--	--	--	--	--
Primary	45.2	56.2	26.5	29.6	18.7	26.6
Secondary	44.6	56.0	27.2	31.3	17.4	24.7
Higher	48.3	50.3	27.7	26.1	20.6	24.2
Wealth Quintile						
Lowest	50.7	57.9	32.8	37.2	17.9	20.7
Lower	44.3	53.4	26.7	28.2	17.5	25.1
Middle	44.3	50.1	26.0	27.4	18.3	22.7
Higher	47.1	59.0	28.6	31.0	18.5	28.1
Highest	41.2	54.3	23.7	28.3	17.6	26.1

1 Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

2 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Contraception Method Mix

Percent distribution of current contraceptive users age 15 to 49, by method type, marital status, and background characteristics.

Background Characteristics	All Users (N = 1046)										Married Users[1] (N = 542)									
	Sterilization	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtnl	Steriliztn	Implant	IUD	DMPA IM	DMPA SC	Pill	EC	Condom	Other Modern	Other Tradtnl
Total																				
All	0.4	15.1	0.5	6.6	1.3	1.6	18.8	14.9	1.0	39.7	0.7	20.3	1.0	8.1	1.9	1.7	9.5	11.0	0.9	44.8
Age Group																				
15-19	0.0	6.4	0.0	2.2	0.8	0.0	40.3	19.6	2.4	28.3	--	--	--	--	--	--	--	--	--	--
20-24	0.0	13.5	0.0	7.5	1.6	1.3	24.4	16.4	2.1	33.3	0.0	17.7	0.0	10.2	3.4	0.0	11.6	13.3	4.0	39.8
25-29	0.0	16.3	0.0	6.2	1.0	2.1	20.5	16.0	0.1	37.8	0.0	23.0	0.0	10.0	1.5	2.0	11.3	13.8	0.3	38.2
30-34	0.6	21.9	1.8	10.8	2.8	3.1	11.7	12.2	0.4	34.8	0.8	23.1	2.4	13.2	3.3	2.9	12.9	8.3	0.5	32.6
35-39	0.3	17.3	1.1	7.4	0.6	1.2	9.6	13.7	0.6	48.3	0.3	21.1	1.4	5.4	0.7	1.5	6.1	10.8	0.7	51.9
40-44	1.9	12.5	1.0	3.4	1.2	1.9	8.0	12.6	0.0	57.6	2.2	13.6	1.1	2.3	1.4	2.2	6.8	11.9	0.0	58.5
45-49	*1.6*	*19.8*	*0.0*	*6.5*	*0.0*	*0.0*	*0.7*	*6.6*	*0.0*	*65.0*	*1.8*	*22.4*	*0.0*	*7.3*	*0.0*	*0.0*	*0.8*	*7.4*	*0.0*	*60.3*
Marital Status																				
Married, or in union	0.7	20.3	1.0	8.1	1.9	1.7	9.5	11.0	0.9	44.8	na	na	na	na	na	na	na	na	na	na
Unmarried, sexually active	0.0	9.1	0.0	5.8	1.0	2.0	25.7	19.4	1.0	36.0	na	na	na	na	na	na	na	na	na	na
Parity																				
0-1 children	0.0	10.8	0.0	9.7	1.2	1.4	19.4	13.6	1.7	42.1	0.0	8.3	0.0	7.2	1.3	0.0	12.1	13.0	2.9	55.3
2-3 children	0.9	23.6	0.7	7.1	2.2	1.7	11.0	12.5	0.9	39.5	1.1	20.5	0.9	7.0	2.7	1.4	8.5	14.7	0.9	42.2
4+ children	0.6	25.3	1.3	8.8	1.5	2.4	7.3	7.3	0.4	45.1	0.7	25.3	1.5	8.9	1.5	2.6	7.0	7.3	0.4	44.9
Education																				
Never	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--	--
Primary	0.0	20.1	0.0	10.5	3.3	2.9	13.5	6.3	2.2	41.3	0.0	20.7	0.0	9.9	3.9	3.4	10.6	2.8	1.2	47.4
Secondary	0.4	16.8	0.7	7.6	1.3	1.8	17.7	13.9	0.9	39.0	0.7	21.4	1.3	8.7	1.8	1.9	6.9	12.4	0.9	44.1
Higher	0.6	5.6	0.0	1.9	0.7	0.5	25.4	21.8	0.9	42.6	1.3	13.0	0.0	4.3	1.3	0.0	21.0	10.1	0.7	48.1
Wealth Quintile																				
Lowest	0.2	27.7	0.8	10.0	3.8	2.0	14.4	4.9	0.9	35.3	0.3	31.7	1.4	12.7	4.3	2.6	6.7	4.6	0.0	35.8
Lower	1.0	15.4	0.0	10.2	1.6	1.8	16.7	13.0	0.7	39.6	2.0	19.9	0.0	12.7	3.0	2.5	3.0	9.1	0.6	47.1
Middle	0.2	11.5	1.0	4.1	1.0	3.7	19.6	16.4	1.3	41.3	0.3	18.6	2.1	1.5	1.5	3.0	13.7	11.4	2.6	45.3
Higher	0.0	12.3	0.0	5.4	0.0	0.3	20.2	20.9	1.6	39.4	0.0	16.2	0.0	7.1	0.0	0.7	10.9	15.8	1.7	47.5
Highest	0.6	9.9	0.9	3.8	0.6	0.6	22.8	17.9	0.4	42.6	1.0	15.7	1.6	6.1	0.9	0.2	12.9	13.6	0.0	48.0

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Modern contraceptive methods

Percentage of women ages 15 to 49 using short-acting and long-acting contraceptive methods by background characteristics.

Background Characteristics	Short Acting Methods [1]		Long Acting Methods [2]	
	All Women (N = 2369)	Married Women [3] (N = 1044)	All Women (N = 2369)	Married Women [3] (N = 1044)
Total				
All	20.0	18.2	7.3	12.1
Age Group				
15-19	14.6	--	1.4	--
20-24	29.7	24.3	7.5	10.1
25-29	27.8	23.8	9.8	14.1
30-34	21.1	22.8	12.5	14.5
35-39	18.3	16.2	10.3	14.6
40-44	13.1	12.7	7.4	8.7
45-49	3.5	4.9	5.4	7.7
Marital Status				
Married, or in union	18.2	na	12.1	na
Unmarried, sexually active	43.2	na	7.2	na
Parity				
0-1 children	25.0	17.5	5.8	4.0
2-3 children	19.7	19.9	13.9	12.7
4+ children	15.4	17.0	15.1	16.7
Education				
Never	--	--	--	--
Primary	17.4	17.9	9.1	11.7
Secondary	19.3	18.3	8.0	13.1
Higher	24.7	18.8	3.0	7.2
Wealth Quintile				
Lowest	18.3	17.8	14.5	19.3
Lower	19.5	16.5	7.3	11.7
Middle	20.4	16.9	5.6	10.5
Higher	22.8	21.4	5.8	9.6
Highest	19.0	18.3	4.7	10.0

1 Short-acting methods include injectables, pills, emergency contraception, condom (male and female), standard days/cycle beads, and LAM.

2 Long-acting methods include IUD, implants, and sterilization (male and female).

3 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Unmet Need for Family Planning

Percent of women ages 15 to 49 with unmet need for family planning (for spacing or limiting), by marital status and background characteristics.

Background Characteristics	All Women (N = 2369)			Married Women[1] (N = 1044)		
	For Spacing	For Limiting	Total	For Spacing	For Limiting	Total
Total						
All	9.1	2.3	11.4	13.0	4.6	17.6
Age Group						
15-19	6.5	0.1	6.6	--	--	--
20-24	11.7	0.6	12.3	24.0	0.4	24.4
25-29	12.0	1.7	13.7	16.4	3.0	19.3
30-34	13.8	1.0	14.8	17.7	1.5	19.2
35-39	10.4	4.5	14.9	11.9	6.3	18.1
40-44	2.4	7.9	10.3	2.9	9.5	12.5
45-49	2.7	6.4	9.1	3.9	7.6	11.4
Marital Status						
Married, or in union	13.0	4.6	17.6	na	na	na
Unmarried, sexually active	13.4	0.7	14.1	na	na	na
Parity						
0-1 children	12.9	0.5	13.4	14.6	0.0	14.6
2-3 children	13.7	1.8	15.4	15.0	2.3	17.3
4+ children	9.4	8.6	18.1	10.6	9.6	20.2
Education						
Never	--	--	--	--	--	--
Primary	16.3	8.7	25.0	17.1	10.9	28.0
Secondary	9.5	2.2	11.7	13.1	4.7	17.7
Higher	5.5	0.1	5.6	10.9	0.3	11.2
Wealth Quintile						
Lowest	14.5	5.3	19.8	16.1	8.8	24.9
Lower	11.7	2.6	14.3	14.5	5.8	20.3
Middle	12.2	1.8	14.0	19.3	4.1	23.4
Higher	5.7	0.9	6.6	10.0	1.5	11.5
Highest	3.7	1.4	5.1	6.5	3.2	9.7

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Need and Demand for Family Planning

Percent of women age 15 to 49 with met need, unmet need, and total demand for family planning, and percentage of demand for family planning satisfied with modern methods, by marital status and background characteristics.

Background Characteristics	All Women (N=2369)					Married Women [2] (N=1044)				
	Any Method	Modern[1] Method	Besoins Unmet Need	Total Demand	Demand Satisfied By Modern	Any Method	Modern[1] Method	Besoins Unmet Need	Total Demand	Demand Satisfied By Modern
Total										
All	45.3	27.3	11.4	56.7	48.2	54.9	30.3	17.6	72.6	41.8
Age Group										
15-19	22.4	16.0	6.6	29.0	55.3	--	--	--	--	--
20-24	55.8	37.2	12.3	68.1	54.6	57.1	34.4	24.4	81.5	42.2
25-29	60.5	37.7	13.7	74.2	50.7	61.4	38.0	19.3	80.7	47.0
30-34	51.5	33.6	14.8	66.3	50.6	55.4	37.3	19.2	74.5	50.1
35-39	55.2	28.5	14.9	70.1	40.7	64.1	30.8	18.1	82.2	37.5
40-44	48.3	20.5	10.3	58.6	35.0	51.7	21.4	12.5	64.1	33.4
45-49	25.2	8.8	9.1	34.3	25.8	31.8	12.6	11.4	43.2	*29.2*
Marital Status										
Married, or in union	54.9	30.3	17.6	72.6	41.8	na	na	na	na	na
Unmarried, sexually active	78.8	50.4	14.1	92.8	54.3	na	na	na	na	na
Parity										
0-1 children	53.1	30.8	13.4	66.5	46.3	48.1	21.5	14.6	62.7	34.3
2-3 children	55.5	33.6	15.4	71.0	47.3	56.5	32.6	17.3	73.8	44.2
4+ children	55.5	30.5	18.1	73.6	41.4	61.2	33.7	20.2	81.4	41.4
Education										
Never	--	--	--	--	--	--	--	--	--	--
Primary	45.2	26.5	25.0	70.2	37.8	56.2	29.6	28.0	84.3	35.1
Secondary	44.6	27.2	11.7	56.3	48.4	56.0	31.3	17.7	73.7	42.5
Higher	48.3	27.7	5.6	53.8	51.4	50.3	26.1	11.2	61.5	42.4
Wealth Quintile										
Lowest	50.7	32.8	19.8	70.6	46.5	57.9	37.2	24.9	82.7	44.9
Lower	44.3	26.7	14.3	58.6	45.6	53.4	28.2	20.3	73.6	38.4
Middle	44.3	26.0	14.0	58.4	44.6	50.1	27.4	23.4	73.5	37.3
Higher	47.1	28.6	6.6	53.7	53.2	59.0	31.0	11.5	70.6	43.9
Highest	41.2	23.7	5.1	46.4	51.1	54.3	28.3	9.7	64.0	44.1

1   Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

2   Married includes married women and those who are living with a man

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Pregnancy Intentions

Unintended births among women age 15 to 49 with one or more births in the previous 5 years or who are currently pregnant: Percent distribution reporting last or current pregnancy was wanted then, wanted later or not wanted at all, by marital status and background characteristics.

Background Characteristics	All Women with 1 or More Births (N=837)			Married[1] Women with 1 or More Births (N=632)		
	Wanted Then	Wanted Later	Not At All	Wanted Then	Wanted Later	Not At All
Total						
All	38.7	54.4	6.9	45.0	47.5	7.5
Age Group						
15-19	*2.6*	*89.1*	*8.3*	--	--	--
20-24	21.5	76.4	2.1	30.6	68.3	1.1
25-29	40.9	57.0	2.0	44.9	53.0	2.2
30-34	48.9	46.4	4.7	50.2	45.1	4.7
35-39	49.9	37.5	12.6	52.2	34.6	13.2
40-44	47.5	38.6	14.0	46.6	40.0	13.4
45-49	--	--	--	--	--	--
Marital Status						
Married, or in union	45.0	47.5	7.5	na	na	na
Unmarried, sexually active	20.0	76.4	3.5	na	na	na
Parity						
0-1 children	30.9	66.4	2.7	48.1	50.8	1.1
2-3 children	47.7	50.3	2.1	53.7	44.8	1.6
4+ children	33.5	50.1	16.4	33.9	49.1	16.9
Education						
Never	--	--	--	--	--	--
Primary	29.8	49.7	20.5	31.8	48.0	20.2
Secondary	35.8	59.1	5.1	42.4	51.9	5.7
Higher	63.6	31.0	5.4	69.9	24.7	5.4
Wealth Quintile						
Lowest	24.7	65.7	9.6	31.1	57.8	11.1
Lower	30.1	63.4	6.6	34.0	58.1	7.9
Middle	36.9	56.8	6.3	42.4	50.5	7.1
Higher	45.3	49.6	5.2	48.3	45.9	5.8
Highest	64.3	29.6	6.0	72.8	22.1	5.1

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Contraceptive Choice

Percent distribution of women age 15 to 49 who are currently using modern contraception[1] reporting on who decided on the contraceptive method, by background characteristics.

Background Characteristics	Self	Jointly with Partner/Provider	Provider/Partner Alone	(N = 547)
Total				
All	48.4	38.1	13.6	
Age Group				
15-19	42.9	29.1	28.0	
20-24	62.3	28.8	8.9	
25-29	49.2	37.6	13.3	
30-34	51.6	39.0	9.4	
35-39	25.9	60.9	13.2	
40-44	*37.3*	*45.5*	*17.2*	
45-49	--	--	--	
Marital Status				
Married, or in union	42.1	48.0	9.9	
Unmarried, sexually active	57.3	26.8	15.8	
Parity				
0-1 children	59.9	25.9	14.2	
2-3 children	47.4	41.8	10.9	
4+ children	47.8	45.1	7.1	
Education				
Never	--	--	--	
Primary	*67.3*	*28.7*	*4.0*	
Secondary	48.9	37.3	13.9	
Higher	38.3	45.6	16.1	
Wealth Quintile				
Lowest	59.7	35.7	4.6	
Lower	52.0	35.1	12.9	
Middle	46.6	33.6	19.8	
Higher	45.3	41.8	12.9	
Highest	36.9	43.8	19.2	

1   Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Method Information Index Components

Percentage of current modern contraceptive [1] users who reported whether provider informed them about other methods, side effects and, if informed of side effects, what to do, by marital status and background characteristics.

Background Characteristics	Informed about other methods [2]		Informed about side effects [2]		Told what to do if experienced side effects [4]		Told about switching methods in future [2]	
	All Users	Married [3] Users	All Users	Married [3] Users	All Users	Married [3] Users	All Users	Married [3] Users
Total	**(547)**	**(278)**	**(547)**	**(278)**	**(211)**	**(142)**	**(547)**	**(278)**
All	31.2	40.0	40.3	52.6	86.7	88.8	36.8	44.9
Age Group								
15-19	9.6	--	16.9	--	--	--	19.3	--
20-24	29.0	*38.1*	37.2	*51.9*	*85.6*	--	33.3	*43.1*
25-29	36.5	45.6	41.7	57.0	*89.7*	*92.6*	41.2	50.3
30-34	38.1	38.7	48.9	46.8	*85.4*	*84.1*	37.8	38.3
35-39	36.0	46.2	49.4	57.9	*91.1*	*90.1*	46.7	48.2
40-44	*27.8*	*26.8*	*41.9*	*42.5*	--	--	*39.8*	*40.2*
45-49	--	--	--	--	--	--	--	--
Marital Status								
Married, or in union	40.0	na	52.6	na	88.8	na	44.9	na
Unmarried, sexually active	19.5	na	25.8	na	82.6	na	27.2	na
Parity								
0-1 children	25.8	*13.4*	36.3	*35.0*	*83.2*	--	32.2	*24.9*
2-3 children	44.0	40.9	55.7	51.6	94.0	92.4	49.0	45.9
4+ children	45.5	45.1	57.7	57.8	86.6	86.2	47.9	48.5
Education								
Never	--	--	--	--	--	--	--	--
Primary	*18.6*	*24.0*	*47.3*	*56.6*	--	--	*37.3*	*45.7*
Secondary	33.9	41.3	42.5	53.2	86.4	87.5	37.4	43.2
Higher	22.9	*44.5*	24.8	*41.5*	--	--	33.9	*53.6*
Wealth Quintile								
Lowest	42.3	37.6	56.9	59.5	89.4	*88.1*	44.2	41.9
Lower	29.5	35.0	41.1	54.5	*82.4*	*82.0*	36.0	43.4
Middle	30.5	*48.7*	34.8	*49.3*	*83.1*	--	34.4	*45.0*
Higher	23.3	*33.4*	30.4	*45.6*	*83.7*	--	28.0	*41.4*
Highest	29.7	*46.6*	36.6	*51.1*	*92.5*	--	41.1	*52.8*

1   Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method
2   Asked of all current/recent modern contraceptive users
3   Married includes married women and those who are living with a man
4   Asked of all current/recent modern contraceptive users who were informed about side effects

na   Not applicable
--   Less than 25 unweighted observations
<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Quality of Care

Percentage of current modern[1] contraceptive users who would return, would refer a friend/relative, or would return and refer a friend/relative to provider, by background characteristics.

Background Characteristics	Percent who would return to provider	Percent who would refer relative/friend	Percent who would return and refer friend/relative
Total	**(N = 528)**	**(N = 528)**	**(N = 528)**
All	64.3	62.3	48.6
Age Group			
15-19	60.5	57.1	48.2
20-24	60.7	57.5	43.8
25-29	64.5	65.1	48.7
30-34	73.3	62.8	51.8
35-39	70.7	74.1	58.2
40-44	*45.1*	*52.0*	*34.5*
45-49	--	--	--
Marital Status			
Married, or in union	68.8	66.3	51.7
Unmarried, sexually active	55.4	54.7	39.8
Parity			
0-1 children	56.7	57.6	40.8
2-3 children	65.9	64.6	48.9
4+ children	73.1	75.8	58.8
Education			
Never	--	--	--
Primary	*70.0*	*84.0*	*63.7*
Secondary	63.1	61.7	47.0
Higher	68.0	56.0	51.5
Wealth Quintile			
Lowest	61.5	76.2	48.4
Lower	63.6	61.0	45.1
Middle	66.4	59.2	49.5
Higher	62.4	50.7	40.5
Highest	68.5	63.2	59.8

1   Modern methods include female and male sterilization, implants, IUD, injectables, pill, emergency contraception, male and female condoms, standard days method, and lactational amenorrhea method

na   Not applicable

--   Less than 25 unweighted observations

<i>   Italics indicate estimate was generated from 25 to 49 unweighted observations

Knowledge of Family Planning

Percentage of women age 15 to 49 who reported receiving family planning information from a health worker in the past 12 months, by marital status and background characteristics.

Background Characteristics	Visited by health worker who discussed family planning		Visited facility and provider discussed family planning		Either visited by health worker or visited facility and discussed family planning	
	All Women	Married[1] Women	All Women	Married[1] Women	All Women	Married[1] Women
Total	**(2369)**	**(1044)**	**(2369)**	**(1044)**	**(2369)**	**(1044)**
All	6.1	5.9	12.1	20.7	17.3	25.3
Age Group						
15-19	5.9	--	4.3	--	9.6	--
20-24	5.2	4.5	16.1	40.4	20.3	44.1
25-29	8.1	7.2	19.5	30.1	26.1	35.8
30-34	6.7	6.2	17.9	22.4	22.3	26.0
35-39	5.5	4.9	13.6	17.4	18.0	20.7
40-44	7.2	7.5	7.9	8.9	15.1	16.3
45-49	3.7	3.9	3.5	4.3	7.1	7.9
Marital Status						
Married, or in union	5.9	na	20.7	na	25.3	na
Unmarried, sexually active	7.0	na	9.1	na	15.1	na
Parity						
0-1 children	6.1	6.6	19.0	26.2	23.5	31.3
2-3 children	6.4	6.1	20.9	24.3	26.3	29.3
4+ children	5.8	6.2	15.5	16.4	20.3	21.3
Education						
Never	--	--	--	--	--	--
Primary	5.2	5.5	18.7	25.4	22.7	29.3
Secondary	6.0	5.6	12.5	20.5	17.6	25.3
Higher	7.2	7.9	8.5	18.5	14.2	23.2
Wealth Quintile						
Lowest	8.6	8.1	17.9	26.9	25.3	34.0
Lower	8.0	9.0	12.5	21.4	19.4	28.1
Middle	5.7	5.6	13.5	22.0	18.3	26.6
Higher	5.7	4.0	9.5	15.5	13.7	18.6
Highest	3.5	3.3	9.2	18.3	12.2	20.4

1 Married includes married women and those who are living with a man

na Not applicable

-- Less than 25 unweighted observations

<i> Italics indicate estimate was generated from 25 to 49 unweighted observations

Response Rates

Response rates of households and individuals, by residence.

Result	Urban	Rural	Total
Household Interviews			
Households Selected	N/A	N/A	2072
Households Occupied	N/A	N/A	1962
Households Interviewed	N/A	N/A	1866
Household Response Rate [1]	0.00%	0.00%	95.11%
Interviews with Women			
Number of Eligible Women [2]	N/A	N/A	2547
Number of Eligible Women Interviewed	N/A	N/A	2369
Eligible Woman Response Rate [3]	0.00%	0.00%	93.01%

1   Household response rate = households interviewed/households selected

2   Eligible women are women ages 15-49 identified in completed household interviews

3   Eligible response rate = eligible women interviewed/eligible women

Sample Error Estimates

Variable	All Women				Married Women [1]			
	Mean	Std Error	CI Lower	CI Upper	Mean	Std Error	CI Lower	CI Upper
Currently using a modern method	0.273	0.017	0.240	0.309	0.303	0.021	0.263	0.347
Currently using a traditional method	0.180	0.012	0.156	0.206	0.246	0.019	0.211	0.285
Currently using any contraceptive method	0.453	0.021	0.411	0.494	0.549	0.020	0.510	0.588
Currently using DMPA-IM	0.066	0.014	0.043	0.100	0.081	0.015	0.056	0.116
Currently using DMPA-SC	0.013	0.004	0.008	0.023	0.019	0.006	0.011	0.033
Currently using condoms	0.149	0.017	0.119	0.185	0.110	0.015	0.083	0.144
Currenlty using implants	0.151	0.023	0.111	0.203	0.203	0.026	0.156	0.261
Chose method by self or jointly in past 12 months	0.864	0.018	0.825	0.896	0.901	0.019	0.857	0.933
Informed by provider about other methods	0.312	0.039	0.240	0.394	0.400	0.038	0.328	0.477
Informed by provider about side effects	0.403	0.037	0.332	0.478	0.526	0.031	0.463	0.587
Satisfied with provider: Would return and refer friend/relative to provider	0.486	0.049	0.389	0.583	0.517	0.050	0.419	0.615
Visited by health worker who talked about family planning in past 12 months	0.061	0.014	0.039	0.095	0.059	0.015	0.036	0.097

1   Married includes married women and those who are living with a man