

PERFORMANCE MONITORING FOR ACTION

PMA KENYA (WEST POKOT)

Results from Phase 1 baseline survey

November-December 2019

OVERALL KEY FINDINGS



There has been a general decrease in modern contraception use since 2016.



51% of the women reported to have received comprehensive methods information during the FP visit.



39% of the pregnancies were unintended.




84% of the users obtained their current modern method from a public health facility

SECTION 1: CONTRACEPTIVE USE, DYNAMICS, AND DEMAND


MODERN CONTRACEPTIVE PREVALENCE

Percent of women age 15-49 currently using modern contraception (mCPR) by marital status



CONTRACEPTIVE PREVALENCE BY METHOD TYPE

Percent of women age 15-49 currently using contraception by method type (PMA Phase 1 n=820)













MODERN CONTRACEPTIVE PREVALENCE BY COUNTY

Percent of women age 15-49 currently using modern contraception by Kenya county



METHOD USE, UNMET NEED, AND DEMAND SATISFIED BY A MODERN METHOD


Percent of women age 15-49 using contraception by method type, unmet need, and demand satisfied by a modern method (PMA Phase 1 = 820)



Demand satisfied by a modern method is use of modern contraceptive methods divided by the sum of unmet need plus total contraceptive use.

TRENDS IN MODERN CONTRACEPTIVE MIX

Percent distribution of modern contraceptive users age 15-49 by method and year (PMA Phase 1 n=165)



MODERN CONTRACEPTIVE METHOD

MD

Percent distribution of married/in union modern contraceptive users age 15-49 by method



12-MONTH DISCONTINUATION RATE

Among women who started an episode of contraceptive use within the two years preceding the survey, the percent of episodes discontinued within 12 months (n=313 episodes)

11% discontinued to become pregnant **53%** discontinued for other reasons

Reasons for discontinuation:



experienced method failure

other method-

related reasons

were concerned over

side effects or health

wanted a more

effective method

had other fertility

related reasons


other/don't know

Discontinued but switched methods:



KEY FINDINGS FOR SECTION 1: CONTRACEPTIVE USE, **DYNAMICS, AND DEMAND**

- There has been a decrease in the use of modern contraception since 2016.
- ●64% of the time, methods were discontinued within one year of starting. 11% did so to become pregnant and 16% switched to another method.
- Implant is the common used method among the married women (45%).
- 39% of the pregnancies were unintended, with 29% mistimed and 10% not wanted



SECTION 2: QUALITY OF FP SERVICES AND COUNSELING

METHOD INFORMATION INDEX PLUS (MII+)

Percent of women who were told about side effects, what to do about side effects, of other methods, and the possibility of switching methods (n=161)

When you obtained your method were you told by the provider about side effects or problems you might have?

Were you told what to do if you experienced side effects or problems?

Were you told by the provider about methods of FP other than the method you received?

Were you told that you could switch to a different method in the future?



Percent of women who responded "Yes" to all four MII+ questions


49% answered "No" to at least one MII+ auestion

answered "Yes" to all four MII+ auestions



DISCUSSED FP IN THE PAST YEAR WITH PROVIDER/CHW Percent of women who received FP information from a provider or community health worker (CHW), by age 24 28 15-19 20-24 25-49 (n=181) (n=150) (n=489)

Yes No



KEY FINDINGS FOR SECTION 2: QUALITY OF FP SERVICES AND COUNSELING

- •About 51% of all women reported to have received comprehensive FP methods information when they obtained their current contraceptive method but 27% reported that the provider did not tell them about their method advantages and disantvantages
- •At the facility, 100% of the clients reported satisfaction with FP services they received while 69% of the clients reported to have obtained their method of choice
- ●Adolescents are about two times less likely to receive FP information from a health provider or a CHW compared to older women

SECTION 3: PARTNER DYNAMICS

PARTNER INVOLVEMENT IN FP DECISIONS

Percent of women who are currently using modern, female controlled methods and agree with the following statements (n=162)

Does your partner know that you are using this method?

Before you started using this method had you discussed the decision to delay or avoid pregnancy with your partner?



Yes No




 ${\it Modern, female controlled methods Includes all modern methods except male sterilization and male condoms}$


Percent of women who are currently using FP and agree with the following statements (n=188)

Would you say

Percent of women who are not currently using FP and agree with the following statements (n=554)



Would you say that not using FP is mainly your decision?



- Joint decision Mainly respondent
- Mainly partner
- Other

KEY FINDINGS FOR SECTION 3: PARTNER DYNAMICS


- ●19% of women using a modern method that can be concealed, reported that their partners do not know that they are using.
- 31% of the women who are using a contraceptive method reported that it is mainly respodent decision to use contraception.
- ●30% of the decisions not to use and 63% of the decisions to use FP are jointly made with the partner.

SECTION 4: WOMEN AND GIRLS' EMPOWERMENT

AGREEMENT WITH FAMILY PLANNING EMPOWERMENT STATEMENTS Percent of all women who strongly agree to strongly disagree with each statement Exercise of choice (self-efficacy, negotiation) for family planning (n=797)


I feel confident telling my provider what is important when selecting an FP method.


I can decide to switch from one FP method to another if I want to.



Existence of choice (motivational autonomy) for family planning (n=790)

Strongly disagree







Neutral

Agree


Disagree

The Women's and Girls' Empowerment (WGE) Index examines existence of choice, exercise of choice, and achievement of choice domains across pregnancy, family planning, and sex outcomes.

Presented results are only for the existence of choice and exercise of choice domains for family planning.

Scores from the above family planning empowerment statements were summed and divided by number of items (7) for average WGE family planning score across both domains.

Range for the combined WGE family planning score is 1-5, with a score of 5 indicating highest empowerment.




Strongly agree



mCPR and intent to use contraception, by categorical WGE score Percent of women using a modern method of contraception and percent of women who intend to use contraception in the next year by categorical WGE score (n=820) 100 80 60 16 40 16 20 9 35 30 12 0 High Lowest Median Highest (5) (1) or low

mCPR and intent to use contraception, by employment

Percent of women using a modern method of contraception and percent of women who intend to use contraception in the next year by employment status (n=820)




KEY FINDINGS FOR SECTION 4: WOMEN AND GIRLS' EMPOWERMENT


- Women who score higher on the empowerment scale are much more likely to be using a modern contraceptive method or to intend to use in the future.
- Women who are employed are more likely to be using or intending to use modern

SECTION 5: ATTITUDES TOWARDS CONTRACEPTION

Intent to use contraception
 mCPR









KEY FINDINGS FOR SECTION 5: ATTITUDES TOWARDS CONTRACEPTION

- Urban women are more likely to agree/strongly agree to the statement that people who use FP have a better quality of life compared to rural women.
- About 5 in every 10 adolescents believe in the myth that adolescents who use FP are promiscuous

SECTION 6: REPRODUCTIVE TIMELINE





KEY FINDINGS FOR SECTION 6: REPRODUCTIVE TIMELINE


- Rural women enter sexual activity earlier, marry earlier, give birth earlier, but initiate contraception later than urban women
- While about 64% of the young women have had first sex by age 18, 22% are married by that age and just 7% have used a contraceptive



SECTION 7: SERVICE DELIVERY POINTS


TRENDS IN METHOD AVAILABILITY: IUD

Public facilities (PMA Phase 1 n=59)




TRENDS IN METHOD AVAILABILITY: IMPLANT

Public facilities (PMA Phase 1 n=59)




TRENDS IN METHOD AVAILABILITY: INJECTABLES

Public facilities (PMA Phase 1 n=59)




TRENDS IN METHOD AVAILABILITY: MALE CONDOMS

Public facilities (PMA Phase 1 n=59)



TRENDS IN METHOD AVAILABILITY: PILLS

Public facilities (PMA Phase 1 n=59)



- Currently in stock and no stockout in last 3 months
- Currently in stock but stockout in last 3 months
- Currently out of stock
- Not offered




FACILITY READINESS

Percent of facilities that provide implants and have a trained provider and instruments/supplies needed for implant insertion/removal (n=59)



Percent of facilities that provide IUDs and have a trained provider and instruments/supplies needed for IUD insertion/removal (n=15)



FEES FOR SERVICES

Percent of facilities where FP clients have to pay fees to be seen by a provider even if they do not obtain FP





of women obtained their current modern method from a public health (facility (n=162)

KEY FINDINGS FOR SECTION 7: SERVICE DELIVERY POINTS

- 84% of the users reported to have obtained their method from a public health facility
- About 75% public SDPs reported not to be offering IUDs in 2019.
- Only 53% of the facilities that provide IUd have a trained provider and instruments/supplies needed for IUD insertion/removal.

TABLES: CONTRACEPTIVE PREVALENCE AND UNMET NEED

A	LL WC	MEN		CPR				mCPR				Unmet need for family planning			
Data source	Round/ Phase	Data collection	Female sample	CPR%	SE	95%	6 CI	mCPR% SE 95% CI		6 CI	Unmet need (%)	SE	95% CI		
PMA 2020	R5	Nov-Dec 2016	434	24.16	6.50	12.95	40.57	22.85	6.75	11.55	40.21	18.70	4.16	11.28	29.40
PMA 2020	R6	Nov-Dec 2017	502	19.24	7.34	8.07	39.27	18.54	7.07	7.80	37.98	17.63	2.79	12.33	24.58
PMA 2020	R7	Nov-Dec 2018	474	22.81	7.26	10.94	41.57	20.17	6.00	10.21	35.96	19.28	2.77	13.92	26.07
PMA	Phase 1	Nov-Dec 2019	820	22.87	4.12	15.53	32.35	19.09	3.31	13.20	26.81	23.54	3.51	17.10	31.49

WC	MEN IN	UNION	ı	CPR				mCPR				Unmet need for family planning			
Data source	Round/ Phase	Data collection	Female sample	CPR%	SE	95%	95% CI mCPR		SE	95% CI		Unmet need (%)	SE	95% CI	
PMA 2020	R5	Nov-Dec 2016	304	27.77	8.50	13.50	48.64	25.86	8.69	11.77	47.69	22.29	5.83	12.20	37.20
PMA 2020	R6	Nov-Dec 2017	346	19.21	7.83	7.61	40.70	19.21	7.83	7.61	40.70	19.16	4.32	11.46	30.26
PMA 2020	R7	Nov-Dec 2018	340	24.27	8.12	11.15	45.00	21.53	6.75	10.46	39.17	20.84	3.21	14.69	28.70
PMA	Phase 1	Nov-Dec 2019	553	23.97	4.71	15.66	34.85	20.71	3.91	13.82	29.85	26.79	3.74	19.82	35.13

PMA Kenya (West Pokot) collects information on knowledge, practice, and coverage of family planning services in 25 enumeration areas selected using a multi-stage stratified cluster design with urban-rural strata. The results are county-level representative. Data were collected between November and December 2019 from 846 households (98.7% response rate), 820 females age 15-49 (99.0% response rate), 66 facilities (93.0% completion rate), and 52 client exit interviews. For sampling information and full data sets, visit www.pmadata.org/countries/kenya.

PMA uses mobile technology and female resident data collectors to support rapid-turnaround surveys to monitor key family planning and health indicators in Africa and Asia. PMA Kenya is led by the Ministry of Health in collaboration with International Centre for Reproductive Health Kenya (ICRHK), National Council for Population and Development, and Kenya National Bureau of Statistics. Overall direction and support are provided by the Bill & Melinda Gates Institute for Population and Reproductive Health at the Johns Hopkins University and Jhpiego. Funding is provided by the Bill & Melinda Gates Foundation.

